

Effective Field Theories for Charmonium


Antonio Vairo

University of Milano and INFN



Quarkonium Working Group

Motivation



- The system is characterized by **two expansion parameters**: α_s and v .
 - (i) hierarchy of scales (\Rightarrow **factorization/effective field theories**)
 - (ii) some of the scales are **perturbative**.
- For these same reasons, charmonium (and quarkonium) are systems where low energy QCD may be studied in a **systematic** way (e.g. **non-perturbative matrix elements, QCD vacuum, confinement, exotica, ...**)

Summary

1. Effective Field Theories: NRQCD, pNRQCD
2. Spectroscopy
 - 2.1 Charm mass
 - 2.2 Higher resonances: pNRQCD potentials
 - 2.3 New Spectroscopy
3. Annihilations
 - 3.1 Inclusive decays
 - 3.2 Electromagnetic decays
4. Production
 - 4.1 Polarization
5. Prospects of $p\bar{p}$ at Fermilab
6. Conclusion


1. EFTs

Quarkonium Scales

Apart from α_s , another small parameter shows up near **threshold**:

$$E \approx 2m + \frac{p^2}{m} + \dots \quad \text{with } v = \frac{p}{m} \ll 1$$


- The perturbative expansion breaks down when $\alpha_s \sim v$:



$$\alpha_s \left(1 + \frac{\alpha_s}{v} + \dots \right) \approx \frac{1}{E - \left(\frac{p^2}{m} + V \right)}$$

- The system is **non-relativistic**: $p \sim mv$ and $E = \frac{p^2}{m} + V \sim mv^2$.

Quarkonium Scales



The mass scale is perturbative:

$$m_b \simeq 5 \text{ GeV}, m_c \simeq 1.5 \text{ GeV}$$

The system is non-relativistic:

$$\Delta_n E \sim m v^2, \Delta_{fs} E \sim m v^4$$

$$v_b^2 \simeq 0.1, v_c^2 \simeq 0.3$$

Non-relativistic bound states are characterized


by at least *three energy scales*

$$m \gg m v \gg m v^2 \quad v \ll 1$$

Normalized with respect to $\chi_b(1P)$ and $\chi_c(1P)$

Quarkonium Scales


Scales get entangled.



Effective Field Theories for Quarkonium

Whenever a system H , described by \mathcal{L}_{QCD} , is characterized by 2 scales $\Lambda \gg \lambda$, observables may be calculated by expanding one scale with respect to the other.

An *effective field theory* makes the expansion in λ/Λ explicit at the Lagrangian level.



$$\frac{\lambda}{\Lambda} = \frac{mv}{m}$$

$$\frac{\lambda}{\Lambda} = \frac{mv^2}{mv}$$

Charmonium Scales

$$m_c \approx 1.5 \text{ GeV} \gg \Lambda_{\text{QCD}}$$

$$m_c v \approx 0.8 \text{ GeV} > \Lambda_{\text{QCD}} \quad \text{for } J/\psi, \eta_c$$


$$m_c v \sim \Lambda_{\text{QCD}} \quad \text{for all higher resonances}$$

As a consequence:

- annihilation, production, **hard scale processes** happen at a **perturbative scale**;
- the bound state is perturbative (i.e. **Coulombic**) perhaps only for the $J/\psi, \eta_c$;
- for all **other charmonium resonances** the bound state is non-perturbative. It will be described by matrix elements, (confining) potentials to be determined on the **lattice**.

NRQCD

NRQCD is the EFT that follows from QCD when $\Lambda = m$




- The **matching** is **perturbative**.
- The Lagrangian is organized as an expansion in $1/m$ and $\alpha_s(m)$:

$$\mathcal{L}_{\text{NRQCD}} = \sum_n c(\alpha_s(m/\mu)) \times O_n(\mu, \lambda)/m^n$$

Suitable to describe **annihilation** and **production** of quarkonium.

pNRQCD

pNRQCD is the EFT for heavy quarkonium that follows from NRQCD when $\Lambda = \frac{1}{r} \sim mv$



- The Lagrangian is organized as an expansion in $1/m$, r , and $\alpha_s(m)$:

$$\mathcal{L}_{\text{pNRQCD}} = \sum_k \sum_n \frac{1}{m^k} \times c_k(\alpha_s(m/\mu)) \times V(r\mu', r\mu) \times O_n(\mu', \lambda) r^n$$

1. Spectroscopy

Low lying $c\bar{c}$

Low lying $c\bar{c}$ states are assumed to realize the hierarchy: $m \gg 1/r \sim mv \gg \Lambda_{\text{QCD}}$

At $mv \gg \mu \gg mv^2$ the EFT is **weakly coupled pNRQCD**; its degrees of freedom are

- $Q-\bar{Q}$ (singlet and octet): $E \sim \Lambda_{\text{QCD}}, mv^2$; $p \lesssim mv$
- Gluons: $E \sim p \sim \Lambda_{\text{QCD}}, mv^2$

Low lying $c\bar{c}$

Low lying $c\bar{c}$ states are assumed to realize the hierarchy: $m \gg 1/r \sim mv \gg \Lambda_{\text{QCD}}$

At $mv \gg \mu \gg mv^2$ the EFT is **weakly coupled pNRQCD**; its degrees of freedom are

- $Q-\bar{Q}$ (singlet and octet): $E \sim \Lambda_{\text{QCD}}, mv^2$; $p \lesssim mv$
- Gluons: $E \sim p \sim \Lambda_{\text{QCD}}, mv^2$

The J/ψ mass at $\mathcal{O}(m\alpha_s^5)$ is

$$E_{J/\psi} = \langle J/\psi | H_s(\mu) | J/\psi \rangle - i \frac{g^2}{3N_c} \int_0^\infty dt \langle 1S | \mathbf{r} e^{it(E_{J/\psi}^{(0)} - H_o)} \mathbf{r} | 1S \rangle \langle \mathbf{E}(t) \mathbf{E}(0) \rangle(\mu)$$

Low lying $c\bar{c}$

Low lying $c\bar{c}$ states are assumed to realize the hierarchy: $m \gg 1/r \sim mv \gg \Lambda_{\text{QCD}}$
At $mv \gg \mu \gg mv^2$ the EFT is **weakly coupled pNRQCD**; its degrees of freedom are

- $Q-\bar{Q}$ (singlet and octet): $E \sim \Lambda_{\text{QCD}}, mv^2$; $p \lesssim mv$
- Gluons: $E \sim p \sim \Lambda_{\text{QCD}}, mv^2$

The J/ψ mass at $\mathcal{O}(m\alpha_s^5)$ is

$$E_{J/\psi} = \langle J/\psi | H_s(\mu) | J/\psi \rangle - i \frac{g^2}{3N_c} \int_0^\infty dt \langle 1S | \mathbf{r} e^{it(E_{J/\psi}^{(0)} - H_o)} \mathbf{r} | 1S \rangle \langle \mathbf{E}(t) \mathbf{E}(0) \rangle(\mu)$$

Brambilla Pineda Soto Vairo 99

From which it follows

$$\bar{m}_c(m_c) = 1.24 \pm 0.020 \text{ GeV}$$

Brambilla Sumino Vairo 01


Higher resonances

Higher $c\bar{c}$ resonances are better studied on the **lattice**.

- QCD ($ma \ll 1$)
- NRQCD (coarse lattices, $ma \gg 1$, no $a \rightarrow 0$)
- pNRQCD (coarse lattices, no $a \rightarrow 0$)

pNRQCD for higher resonances

- All quarks with energy $\gg mv^2$ and momentum $\gg mv$ are integrated out.
- All gluonic excitations between heavy quarks are integrated out since they develop a gap of order Λ_{QCD} with the static $Q\bar{Q}$ energy.



Bali et al. 98

($r_0 \simeq 0.5$ fm)

⇒ The singlet quarkonium field S of energy mv^2 and momentum mv is the only the degree of freedom of pNRQCD (up to ultrasoft light quarks, e.g. pions).

pNRQCD for higher resonances


- All quarks with energy $\gg mv^2$ and momentum $\gg mv$ are integrated out.
- All gluonic excitations between heavy quarks are integrated out since they develop a gap of order Λ_{QCD} with the static $Q\bar{Q}$ energy.

$$\mathcal{L} = \text{Tr} \left\{ S^\dagger \left(i\partial_0 - \frac{\mathbf{p}^2}{m} - V_s \right) S \right\}$$

- The idea is to calculate once for ever the potentials on the lattice and determine the spectrum by solving the Schrödinger equation.

Static potential


$$V_s^{(0)} = \lim_{T \rightarrow \infty} \frac{i}{T} \ln \langle W(r \times T) \rangle = \lim_{T \rightarrow \infty} \frac{i}{T} \ln \langle \boxed{} \rangle$$



1/m potential

$$V_s^{(1)} = -\frac{1}{2} \int_0^\infty dt t \langle \text{box} \rangle$$


Brambilla Pineda Soto Vairo 00




Koma, Koma and Wittig/QWG 06

Spin-dependent potentials


$$2 \int_0^\infty d\tau \tau \langle\langle B_y(\mathbf{r}, 0) E_z(\mathbf{r}, \tau) \rangle\rangle$$




$$2 \int_0^\infty d\tau \tau \langle\langle B_y(\mathbf{0}, 0) E_z(\mathbf{r}, \tau) \rangle\rangle$$



$$2 \int_0^\infty d\tau [\langle\langle B_x(\mathbf{0}, 0) B_x(\mathbf{r}, \tau) \rangle\rangle - (x \rightarrow y)]$$




$$2 \int_0^\infty d\tau [\langle\langle B_x(\mathbf{0}, 0) B_x(\mathbf{r}, \tau) \rangle\rangle + 2(x \rightarrow y)]$$



States near or above threshold


- In general, for states near or above threshold a systematic treatment does not exist so far. Most of the existing analyses rely on models (e.g. the Cornell coupled channel model).
- However one may still exploit an expansion in α_s and v . In some cases one may develop an EFT owing to special dynamical conditions.
 - A possible exotic (hybrid) is the $Y(4260)$.
 - An example is the $X(3872)$ interpreted as a $D^0 \bar{D}^{*0}$ or $\bar{D}^0 D^{*0}$ molecule. In this case, one may take advantage of the unnaturally (and accidentally) large $D^0 \bar{D}^{*0}$ scattering length.

Braaten Kusunoki 03




$h_c(3523)$

CLEO 05
E835 05




$\eta_c(2S)(3630)$

BaBar 04
CLEO 04
Belle 02




X(3872)

CDF D0/OWG 04
Belle 02
BaBar 05




Z(3930)

Belle 05



X(3940)

Belle 05



Y(4260)


BaBar 05

2. Annihilations

NRQCD factorization

$$\Gamma(H \rightarrow \text{LH}) = \sum_n \frac{2 \text{Im} f^{(n)}}{m^{d_n-4}} \langle H | \psi^\dagger K^{(n)} \chi \chi^\dagger K'^{(n)} \psi | H \rangle$$

$$\Gamma(H \rightarrow \text{EM}) = \sum_n \frac{2 \text{Im} f_{\text{em}}^{(n)}}{m^{d_n-4}} \langle H | \psi^\dagger K^{(n)} \chi | \text{vac} \rangle \langle \text{vac} | \chi^\dagger K'^{(n)} \psi | H \rangle$$



Bodwin et al 95

NRQCD matrix elements

- By fitting charmonium P -wave decay data

$\langle O_1(^1P_1) \rangle_{h_c(1P)} \approx 8.1 \times 10^{-2} \text{ GeV}^5$ and $\langle O_8(^1S_0) \rangle_{h_c(1P)} \approx 5.3 \times 10^{-3} \text{ GeV}^3$
in $\overline{\text{MS}}$ and at the factorization scale of 1.5 GeV.

Maltoni 00

- In quenched lattice simulations

$\langle O_1(^1P_1) \rangle_{h_c(1P)} \approx 8.0 \times 10^{-2} \text{ GeV}^5$, $\langle O_8(^1S_0) \rangle_{h_c(1P)} \approx 4.7 \times 10^{-3} \text{ GeV}^3$ and
 $\langle O_1(^1S_0) \rangle_{\eta_c(1S)} \approx 0.33 \text{ GeV}^3$
in $\overline{\text{MS}}$ and at the factorization scale of 1.3 GeV.

Bodwin Sinclair Kim 96

- In lattice simulations with three light-quark flavors (extrapolation)


$\langle O_1(^1S_0) \rangle_{\eta_b(1S)} \approx 4.1 \text{ GeV}^3$, $\langle O_1(^1P_1) \rangle_{h_b(1P)} \approx 3.3 \text{ GeV}^5$ and
 $\langle O_8(^1S_0) \rangle_{h_b(1P)} \approx 5.9 \times 10^{-3} \text{ GeV}^3$
in $\overline{\text{MS}}$ and at the factorization scale of 4.3 GeV.

Bodwin Sinclair Kim 01

Some further recent (quenched) determinations are in Bodwin Lee Sinclair 05

pNRQCD factorization

$$\langle H | \psi^\dagger K^{(n)} \chi \chi^\dagger K'^{(n)} \psi | H \rangle = |R(0)|^2 \times \int dt t^n \langle G(t) G(0) \rangle$$



P-wave decays at $\mathcal{O}(mv^5)$

- NRQCD

$$\Gamma(\chi_J \rightarrow \text{LH}) = 9 \text{Im } f_1 \frac{|R'(0)|^2}{\pi m^4} + \frac{2 \text{Im } f_8}{m^2} \langle \chi | O_8(^1S_0) | \chi \rangle$$
$$\Gamma(\chi_J \rightarrow \gamma\gamma) = 9 \text{Im } f_{\gamma\gamma} \frac{|R'(0)|^2}{\pi m^4} \quad J = 0, 2$$

* Bottomonium and charmonium P-wave decays depend on 6 non-perturbative parameters.


- pNRQCD

$$\langle \chi | O_8(^1S_0) | \chi \rangle = \frac{|R'(0)|^2}{18\pi m^2} \mathcal{E}; \quad \mathcal{E} \equiv \int_0^\infty dt t^3 \langle \text{Tr}(g\mathbf{E}(t) g\mathbf{E}(0)) \rangle$$

* The quarkonium state dependence factorizes.

* Bottomonium and charmonium P-wave decays depend on 4 non-perturbative parameters.

Bottomonium P -wave decays



$$\frac{\Gamma(\chi_{b0}(1P) \rightarrow \text{LH})}{\Gamma(\chi_{b1}(1P) \rightarrow \text{LH})} = \frac{\Gamma(\chi_{b0}(2P) \rightarrow \text{LH})}{\Gamma(\chi_{b1}(2P) \rightarrow \text{LH})} = 8.0 \pm 1.3$$

$$(\text{CleoIII 02}) = 19.3 \pm 9.8$$

Charmonium P-wave decays

Ratio	PDG04	PDG00	LO	NLO
$\frac{\Gamma(\chi_{c0} \rightarrow \gamma\gamma)}{\Gamma(\chi_{c2} \rightarrow \gamma\gamma)}$	5.1 ± 1.1	13 ± 10	3.75	≈ 5.43
$\frac{\Gamma(\chi_{c2} \rightarrow l.h.) - \Gamma(\chi_{c1} \rightarrow l.h.)}{\Gamma(\chi_{c0} \rightarrow \gamma\gamma)}$	410 ± 100	270 ± 200	≈ 347	≈ 383
$\frac{\Gamma(\chi_{c0} \rightarrow l.h.) - \Gamma(\chi_{c1} \rightarrow l.h.)}{\Gamma(\chi_{c0} \rightarrow \gamma\gamma)}$	3600 ± 700	3500 ± 2500	≈ 1300	≈ 2781
$\frac{\Gamma(\chi_{c0} \rightarrow l.h.) - \Gamma(\chi_{c2} \rightarrow l.h.)}{\Gamma(\chi_{c2} \rightarrow l.h.) - \Gamma(\chi_{c1} \rightarrow l.h.)}$	7.9 ± 1.5	12.1 ± 3.2	2.75	≈ 6.63
$\frac{\Gamma(\chi_{c0} \rightarrow l.h.) - \Gamma(\chi_{c1} \rightarrow l.h.)}{\Gamma(\chi_{c2} \rightarrow l.h.) - \Gamma(\chi_{c1} \rightarrow l.h.)}$	8.9 ± 1.1	13.1 ± 3.3	3.75	≈ 7.63

$$m_c = 1.5 \text{ GeV} \quad \alpha_s(2m_c) = 0.245$$

mainly from E835 (χ_{c0} , total width and $\gamma\gamma$)

also from Belle ($\chi_{c0} \rightarrow \gamma\gamma$) and CLEO, BES

$$\Gamma(\eta_c \rightarrow LH)/\Gamma(\eta_c \rightarrow \gamma\gamma)$$

- Large $\beta_0\alpha_s$ contributions.

$$\frac{\Gamma(\eta_c \rightarrow LH)}{\Gamma(\eta_c \rightarrow \gamma\gamma)} \approx (1.1 \text{ (LO)} + 1.0 \text{ (NLO)}) \times 10^3 = 2.1 \times 10^3$$

$$\frac{\Gamma(\eta_c \rightarrow LH)}{\Gamma(\eta_c \rightarrow \gamma\gamma)} = (3.3 \pm 1.3) \times 10^3 \text{ (EXP)}$$


$$\Gamma(\eta_c \rightarrow LH) / \Gamma(\eta_c \rightarrow \gamma\gamma)$$

- Large $\beta_0 \alpha_s$ contributions.

$$\frac{\Gamma(\eta_c \rightarrow LH)}{\Gamma(\eta_c \rightarrow \gamma\gamma)} \approx (1.1 \text{ (LO)} + 1.0 \text{ (NLO)}) \times 10^3 = 2.1 \times 10^3$$

$$\frac{\Gamma(\eta_c \rightarrow LH)}{\Gamma(\eta_c \rightarrow \gamma\gamma)} = (3.3 \pm 1.3) \times 10^3 \text{ (EXP)}$$


- scheme dependence
- renormalons



$$\frac{\Gamma(\eta_c \rightarrow LH)}{\Gamma(\eta_c \rightarrow \gamma\gamma)} = (3.01 \pm 0.30 \pm 0.34) \times 10^3$$

$$\Gamma(J/\psi \rightarrow e^+e^-)/\Gamma(\eta_c \rightarrow \gamma\gamma)$$

- Large logarithms.



$$\mathcal{R}_c = \frac{\Gamma(J/\psi \rightarrow e^+e^-)}{\Gamma(\eta_c \rightarrow \gamma\gamma)}$$

3. Production

Quarkonium Production

- There is **no** formal proof of the NRQCD **factorization** yet.
- The relevant **4-fermion operators** are


$$\psi^\dagger K^{(n)} \chi a_H^\dagger a_H \chi^\dagger K'^{(n)} \psi$$

Recently it has been proved that the **cancellation of the IR divergences at NNLO** requires the modification of the 4 fermion operators into


$$\psi^\dagger K^{(n)} \chi \phi_l^\dagger(0, \infty) a_H^\dagger a_H \phi_l(0, \infty) \chi^\dagger K'^{(n)} \psi$$
$$\phi_l(0, \infty) = \text{P exp} \left(-ig \int_0^\infty d\lambda l \cdot A(\lambda l) \right), \quad l^2 = 1$$

Charmonium Production at the Tevatron

Octet contributions dominate in production at high p_T .



$$p\bar{p} \rightarrow J/\psi + X$$





$$p\bar{p} \rightarrow \psi(2S) + X$$

Charmonium Polarization at the Tevatron

- For large p_T quarkonium production, gluon fragmentation via the color-octet mechanism dominates: $\langle O_8^{J/\psi}({}^3S_1) \rangle$.
- At large p_T the gluon is nearly on mass shell and so is transversely polarized.
- In color octet gluon fragmentation, most of the gluon's polarization is transferred to the J/ψ .
- Radiative corrections, color singlet production dilute this.
- In the case of the J/ψ feeddown is important:
feeddown from χ_c states is about 30% of the J/ψ sample and dilutes the polarization.
- feeddown from $\psi(2S)$ is about 10% of the J/ψ sample and is largely transversely polarized.
- *Spin-flipping terms are assumed suppressed. But This strictly depends on the **power counting**.
If they are not, polarization may dilute at high p_T .*


Charmonium Polarization at the Tevatron




$$\frac{d\sigma}{d\cos\theta} \propto 1 + \alpha \cos^2\theta$$

$\alpha = 1$ is completely transverse $\alpha = -1$ is completely longitudinal.


Charmonium Polarization at the Tevatron



Charmonium Polarization at the Tevatron



Charmonium Polarization at the Tevatron



5. Prospects of $p\bar{p}$ at Fermilab

E760-E835 legacy

$p\bar{p}$ to charmonium is good for

- precise determination of **known** resonances
 - * mass measurements at about 0.1 MeV level,
 - * total width measurements (ideal for narrow states);
- through the detection of:
EM final states (e.g. electrons and photons) few body final states.

E760-E835 legacy

$p\bar{p}$ to charmonium was limited by

- non hermicity of the detector;
- low energy photon threshold: 20 MeV;
- calorimeter granularity;
- multiple scattering for tracks below 1 GeV;
- physical occupancy of the jet target;
- 2x extra rate induced by $e\text{-}\bar{p}$ interactions;
- no momentum measurement on hadrons (no magnet).

The current generation of B factories will **not** have enough statistics to measure $p\bar{p}$ coupling to newly discovered states. Probably super B factories will be able to measure some in B decays (limited to $J=0,1$). Panda is scheduled to start after 2014, i.e. in the super B era.

The **antiproton source at Fermilab** (the antiproton ring should have originally supplied beam to BTeV) provides (after 2009 and before 2014: data taking from 2011) a potential tool to understand XYZ's. What is needed is:

- an improved detector and a target with respect to the old E835;
- B factories that tell where to look.

Conclusion

Charmonium physics provides a place where low-energy QCD and its rich structure may be studied in a **controlled** and **systematic** way, combining **perturbative QCD**, **lattice calculations**, and **effective field theory** analytical methods.

B factories have immensely boosted charmonium studies in the last years, which makes the contribution to charmonium physics that **$p\bar{p}$ machines** at Fermilab and GSI have provided and could/will provide in the future even more valuable.