



Dissertation Agnes Lundborg
The Charm of Excited Glue
Uppsala, February 23, 2007

Exotic Particles in Meson Spectroscopy (1)

Constituents of matter



Constituents of a proton: Quarks and Gluons



Flavor (q)	Mass [MeV]	Q/e
u	5	+ 2/3
d	7	- 1/3
s	95	- 1/3
c	1250	+ 2/3
b	4200	- 1/3
t	174200	+ 2/3

Theory: Quantum Chromo Dynamics (QCD)

Naive Quark Model:

Baryons: qqq

Proton = uud

Antiproton = $\bar{u}\bar{u}\bar{d}$

Mesons : $q\bar{q}$

$\pi^+ = u\bar{d}$

$\pi^- = \bar{u}d$

Characterization: Mass (m), Decaytime (τ) or Width ($\Gamma = \hbar/\tau$), Decay modes, Quantum numbers (J^{PC})

Exotic Particles in Meson Spectroscopy (2)

Problems with the naive Quark Model








– DIS-Experiments

– Mass of the Hadrons much heavier than expected

Example (Proton): $M \approx 17 \text{ MeV}$
 $M_{\text{exp.}} = 938 \text{ MeV}$ } Large dynamical mass generation process

– Many of the new discoveries don't fit into the picture (see later)

– QCD allows many more states than described by the naive model (**Exotic Particles**)

$(gg), (ggg)$	Glue-Balls			Soliton-Type States		
$(\bar{q}qg)$	Hybrids					
$(qq) (\bar{q}\bar{q})$	Diquonium		} Quark-Molecules	$(qq) (qq\bar{q})$	Penta Quark States	
$(q\bar{q}) (q\bar{q})$	Mesonium			$(qqq) (qq\bar{q})$	Dibaryons	
$(qqq) (\bar{q}\bar{q}\bar{q})$	Baryonium					

New feature: Spin-exotic quantum numbers possible, not allowed in $q\bar{q}$ ($J^{PC} = 0^{+-}, 1^{-+}, \dots$)

Exotic Particles in Meson Spectroscopy (3)

Glueballs (gg)

Predictions:

Masses:

1.5-5.0 GeV/c² (Ground state found? ;
Candidates for further states?)


Quantum numbers:

Several spin exotics (oddballs), e.g.
 $J^{PC} = 2^{+-}$ (4.3 GeV/c²)

Widths: ≥ 100 MeV/c²


- Decay into two lighter glueballs often forbidden because of q.-n.
- No mixing effects for oddballs

Decays: $\phi\phi$, $\phi\eta$, $\eta\pi$




Exotic Particles in Meson Spectroscopy (4)

Charmonium Hybrids ($c\bar{c}g$, H_c)



Decay modes:
 $J/\psi\omega$; $D^*\bar{D}$


Small overlap
with $c\bar{c}$ -states




42] K. Juge, J. Kuti, and C. Morningstar,
Phys. Rev. Lett. 90, 161601 (2003).

Exotic Particles in Meson Spectroscopy (5)

Candidates for Exotics in the low mass sector (mainly from LEAR/CERN)



LEAR-Ring



Exotic Particles in Meson Spectroscopy (6)

Glue-Ball Candidate

$f_0(1500)$ (Best candidat for the Glueball-ground state)


Production : $p\bar{p} \rightarrow f_0(1500)\pi^0$ (Crystal Barrel/LEAR)

Decays : $f_0(1500) \rightarrow 2\pi, 4\pi, \eta\eta, \eta\eta', K\bar{K}$

$M = (1505 \pm 9) \text{ MeV} ; \Gamma = (111 \pm 12) \text{ MeV} ; J^{PC} = 0^{++}$

Exotic?

- Surplus state in 0^{++} -Nonett
- Relatively narrow width
- Decays in particles, which contain u, d and s-Quarks
- Mass and Quantum numbers in good agreement with Lattice QCD-predictions for the Glueball-ground state



Exotic Particles in Meson Spectroscopy (7)

Hybrid Candidates with exotic quantum number combination

$\pi_1(1400) / \pi_1(1600)$

Production/Decays :

$\pi^- p \rightarrow \pi_1(1400) p$ (E835/BNL) and $\bar{p} n \rightarrow \pi_1(1400) \pi^0$ (Crystal Barrel/LEAR)
 $\hookrightarrow \eta \pi^-$ $\hookrightarrow \eta \pi^-$

$\pi^- p \rightarrow \pi_1(1600) p$ (E835 BNL) and $\bar{p} p \rightarrow \pi_1(1600) \pi^+$ (Crystal Barrel/LEAR)
 $\hookrightarrow \pi^- \eta$ $\hookrightarrow \pi^- \eta$



$M \approx 1400, 1600 \text{ MeV}$; $\Gamma \approx 300 \text{ MeV}$; $J^{PC} = 1^{+-}$ (Exotic Q.-N., At variance with naive Quark-model)

Exotic?

Exotic J^{PC} -combination

Hybrids?

Multi-Quark-states?




Exotic Particles in Meson Spectroscopy (8)

Contributions of A. Lundborg:


Look for Glueballs in radiative decays:

$(c\bar{c}) \rightarrow \gamma + f_0(1500), \dots$


Production of $(c\bar{c})$: $e^+e^- \rightarrow (c\bar{c})$



Positronium




Charmonium




Exotic Particles in Meson Spectroscopy (9)

Beijing Electron Positron Collider (BEPC)



$$e^+e^- \rightarrow (c\bar{c})$$

BES II



Exotic Particles in Meson Spectroscopy (10)

Results: $\psi' \rightarrow \gamma + X$

$\hookrightarrow \pi^+ \pi^-, K^+ K^-$

$$\text{BR}(\psi' \rightarrow \gamma f_2(1270) \rightarrow \gamma \pi^+ \pi^-) = (2.2 \pm 0.1_{-0.2}^{+0.2}) \times 10^{-4}$$

$$\text{BR}(\psi' \rightarrow \gamma f_0(1500) \rightarrow \gamma \pi^+ \pi^-) = (1.5 \pm 0.7_{-0.4}^{+0.9}) \times 10^{-5}$$

$$\text{BR}(\psi' \rightarrow \gamma f_0(1710) \rightarrow \gamma \pi^+ \pi^-) = (2.4 \pm 0.6_{-1.1}^{+0.7}) \times 10^{-5}$$



$$\text{BR}(\psi' \rightarrow \gamma f_4(2050) \rightarrow \gamma \pi^+ \pi^-) = (2.8 \pm 0.9_{-0.6}^{+0.8}) \times 10^{-5}$$

$$\text{BR}(\psi' \rightarrow \gamma f_0(2200) \rightarrow \gamma \pi^+ \pi^-) = (4.6 \pm 1.0_{-1.0}^{+4.5}) \times 10^{-5}$$

$$\text{BR}(\psi' \rightarrow \gamma f_2(1270) \rightarrow \gamma K^+ K^-) = (1.9 \pm 0.6_{-0.6}^{+1.0}) \times 10^{-5}$$

$$\text{BR}(\psi' \rightarrow \gamma f_2'(1525) \rightarrow \gamma K^+ K^-) = (6.9 \pm 4.4_{-2.1}^{+4.1}) \times 10^{-6}$$

$$\text{BR}(\psi' \rightarrow \gamma f_0(1710) \rightarrow \gamma K^+ K^-) = (3.1 \pm 0.6_{-0.7}^{+1.1}) \times 10^{-5}$$



Significance much higher than for BES I-data. Further improvements expected from BES III

For the first time BR was given for $\psi' \rightarrow \gamma f_0(1500)$


The appearances of $f_0(1500)$ and $f_0(1710)$ are consistent with the hypothesis, that $f_0(1500)$ is the ground state glue ball

Exotic Particles in Meson Spectroscopy (11)

Discussion of the $J^{PC} = 0^{++}$ nonet

Nine open slots, but twelve candidates

Possible scenario:




$a_0(980), f_0(980)$: 4 quark states

Exotic Particles in Meson Spectroscopy (12)


Candidates for Exotics in the high mass sector (BaBar/BELLE/CLEO-c/...)

$$e^+e^- \rightarrow Y(4s) \rightarrow B\bar{B} \\ \rightarrow (c\bar{c}) + (J/\psi) + X, \dots$$

BaBar



Belle




Exotic Particles in Meson Spectroscopy (13)

Candidates for exotics states


$X(3872)$, $J^{PC} = 1^{++} (?)$, $\Gamma < 2.3$ MeV!!

$\hookrightarrow \pi^+ \pi^- J/\psi$



$Y(3949)$, $J^{PC} = ?$, $\Gamma = (92 \pm 24)$ MeV


$\hookrightarrow \omega J/\psi$



$Y(4260)$, $J^{PC} = 1^{--}$, $\Gamma = (88 \pm 23)$ MeV

$\hookrightarrow \pi^+ \pi^- J/\psi$



$H_c?$



Exotic Particles in Meson Spectroscopy (14)


Contributions of A. Lundborg

High energies antiprotons at GSI: FAIR-Project



Exotic Particles in Meson Spectroscopy (15)

The PANDA-Detector



Detector requirements

- full angular acceptance and angular resolution for charged particles and γ , π^0
- particle identification (π , K , e , μ) in the range up to ~ 8 GeV/c
- high momentum resolution in a wide energy range
- high rate capabilities, especially in interaction point region and forward detector :
expected interaction rate $\sim 10^7/s$
- precise vertex reconstruction for fast decaying particles

Exotic Particles in Meson Spectroscopy (16)

Simulation of a Charmed Hybrid (H_c)

$\bar{p}p \rightarrow H_c \pi^0 / \eta$; $H_c \rightarrow \chi_{c_1} (\pi^0 \pi^0)_{S\text{-wave}}$; $\chi_{c_1} \rightarrow \gamma J/\psi$; $J/\psi \rightarrow e^+ e^- (\mu^+ \mu^-) \rightarrow \ell^+ \ell^- + 7\gamma$'s
Assumption: $M(H_c) = 4.28 \text{ GeV}$ $\Gamma(H_c) = 20 \text{ MeV}$ $\sigma(\bar{p}p \rightarrow H_c \pi^0) = 100 \text{ pb}$


Problem: Signal/Hadronic background $\approx 10^{-9}$

Good test case for requests of EMC-properties: Energy resolution/Energy threshold/
solid angle coverage

Work not yet finished because of lacking description of components and kinematical fits

Preliminary, very encouraging results:

- Channel measurable in spite of a high hadronic background



- Many detailed results relevant for the final design of EMC

Exotic Particles in Meson Spectroscopy (17)


Simulation of $\gamma\gamma$ -decay of the Charmonium ground state: $\eta_c \rightarrow \gamma\gamma$

Reasons for a refined measurement:

Precision determination of $\alpha_s(2.9 \text{ GeV})$

Precision determination of total width

Result:



Peak/Background: $5.4 \pm 0.4 \pm 1.1$

(In spite of dominant background channels: $\bar{p}p \rightarrow \pi^0\pi^0, \pi^0\gamma$)

Exotic Particles in Meson Spectroscopy (18)

Production of $(c\bar{c})$ -states in $p\bar{p}$ -annihilations (Lundborg, Barnes, Wiedner)

Example: $p\bar{p} \rightarrow J/\psi + \pi^0(m)$

Knowledge of cross section very important for PANDA-research program

Until now: Only one measurement, only one calculation

New Method: Use $J/\psi \rightarrow m\bar{p}p$ branching ratio to determine the cross section

Good result for $J/\psi + \pi^0$, in reasonable agreement with data

Resumee

- The thesis of A. Lundborg deals with a very exciting part of modern Hadron Physics
- A. Lundborg has used high level and very sophisticated tools for her analysis
- The results contributed essentially to a further understanding of exotic hadronic states